

Latihan Soal Rekursi

7. Give a recursive definition of the sequence $\{a_n\}$, $n = 1, 2, 3, \dots$ if
- | | |
|------------------|--------------------|
| a) $a_n = 6n.$ | b) $a_n = 2n + 1.$ |
| c) $a_n = 10^n.$ | d) $a_n = 5.$ |

A string that contains only 0s, 1s, and 2s is called a **ternary string**.

13. a) Find a recurrence relation for the number of ternary strings of length n that do not contain two consecutive 0s.
- b) What are the initial conditions?
- c) How many ternary strings of length six do not contain two consecutive 0s?

1. Determine which of these are linear homogeneous recurrence relations with constant coefficients. Also, find the degree of those that are.

a) $a_n = 3a_{n-1} + 4a_{n-2} + 5a_{n-3}$

b) $a_n = 2na_{n-1} + a_{n-2}$ c) $a_n = a_{n-1} + a_{n-4}$

d) $a_n = a_{n-1} + 2$ e) $a_n = a_{n-1}^2 + a_{n-2}$

f) $a_n = a_{n-2}$ g) $a_n = a_{n-1} + n$

3. Solve these recurrence relations together with the initial conditions given.

a) $a_n = 2a_{n-1}$ for $n \geq 1$, $a_0 = 3$

b) $a_n = a_{n-1}$ for $n \geq 1$, $a_0 = 2$

c) $a_n = 5a_{n-1} - 6a_{n-2}$ for $n \geq 2$, $a_0 = 1$, $a_1 = 0$

d) $a_n = 4a_{n-1} - 4a_{n-2}$ for $n \geq 2$, $a_0 = 6$, $a_1 = 8$

e) $a_n = -4a_{n-1} - 4a_{n-2}$ for $n \geq 2$, $a_0 = 0$, $a_1 = 1$

f) $a_n = 4a_{n-2}$ for $n \geq 2$, $a_0 = 0$, $a_1 = 4$

g) $a_n = a_{n-2}/4$ for $n \geq 2$, $a_0 = 1$, $a_1 = 0$

EXAMPLE 11 Find all solutions of the recurrence relation

Extra
Examples



$$a_n = 5a_{n-1} - 6a_{n-2} + 7^n.$$

Solution: This is a linear nonhomogeneous recurrence relation. The solutions of its associated homogeneous recurrence relation

$$a_n = 5a_{n-1} - 6a_{n-2}$$

are $a_n^{(h)} = \alpha_1 \cdot 3^n + \alpha_2 \cdot 2^n$, where α_1 and α_2 are constants. Because $F(n) = 7^n$, a reasonable trial solution is $a_n^{(p)} = C \cdot 7^n$, where C is a constant. Substituting the terms of this sequence into the recurrence relation implies that $C \cdot 7^n = 5C \cdot 7^{n-1} - 6C \cdot 7^{n-2} + 7^n$. Factoring out 7^{n-2} , this equation becomes $49C = 35C - 6C + 49$, which implies that $20C = 49$, or that $C = 49/20$. Hence, $a_n^{(p)} = (49/20)7^n$ is a particular solution. By Theorem 5, all solutions are of the form

$$a_n = \alpha_1 \cdot 3^n + \alpha_2 \cdot 2^n + (49/20)7^n.$$



23. Consider the nonhomogeneous linear recurrence relation
 $a_n = 3a_{n-1} + 2^n$.

- a)** Show that $a_n = -2^{n+1}$ is a solution of this recurrence relation.
- b)** Use Theorem 5 to find all solutions of this recurrence relation.
- c)** Find the solution with $a_0 = 1$.

25. a) Determine values of the constants A and B such that
 $a_n = An + B$ is a solution of recurrence relation $a_n = 2a_{n-1} + n + 5$.

- b)** Use Theorem 5 to find all solutions of this recurrence relation.
- c)** Find the solution of this recurrence relation with $a_0 = 4$.