# Latihan Soal Lanjutan Relasi

**3.** List the ordered pairs in the relations on {1, 2, 3} corresponding to these matrices (where the rows and columns correspond to the integers listed in increasing order).


a) 
$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$
b) 
$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{bmatrix}$$
c) 
$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

Determine whether the relations represented by the matrices in Exercise 3 are reflexive, irreflexive, symmetric, antisymmetric, and/or transitive.


For reflexivity we want all 1's on the main diagonal; for irreflexivity we want all 0's on the main diagonal; for symmetry, we want the matrix to be symmetric about the main diagonal (equivalently, the matrix equals its transpose); for antisymmetry we want there never to be two 1's symmetrically placed about the main diagonal (equivalently, the meet of the matrix and its transpose has no 1's off the main diagonal); and for transitivity we want the Boolean square of the matrix (the Boolean product of the matrix and itself) to be "less than or equal to" the original matrix in the sense that there is a 1 in the original matrix at every location where there is a 1 in the Boolean square.

In Exercises 23–28 list the ordered pairs in the relations represented by the directed graphs.






25.




27.




Recall that the relation is reflexive if there is a loop at each vertex; irreflexive if there are no loops at all; symmetric if edges appear only in **antiparallel** pairs (edges from one vertex to a second vertex and from the second back to the first); antisymmetric if there is no pair of antiparallel edges; asymmetric if is both antisymmetric and irreflexive; and transitive if all paths of length 2 (a pair of edges (x, y) and (y, z)) are accompanied by the corresponding path of length 1 (the edge (x, z)). The relation drawn in Exercise 26 is

23.




**31.** Determine whether the relations represented by the directed graphs shown in Exercises 23–25 are reflexive, irreflexive, symmetric, antisymmetric, and/or transitive.

25.




27.




- **1.** Let *R* be the relation on the set {0, 1, 2, 3} containing the ordered pairs (0, 1), (1, 1), (1, 2), (2, 0), (2, 2), and (3, 0). Find the
  - a) reflexive closure of R. b) symmetric closure of R.


5.



6.




7.



Symmetric closure? Reflexive closure?


**16.** Determine whether these sequences of vertices are paths in this directed graph.



**b**) *b*, *e*, *c*, *b*, *e* 

**d**) *b*, *c*, *e*, *d*, *a*, *a*, *b* 

**f**) a, a, b, b, c, c, b, e, d



**17.** Find all circuits of length three in the directed graph in Exercise 16.

**21.** Let *R* be the relation on the set of all students containing the ordered pair (a, b) if a and b are in at least one common class and  $a \neq b$ . When is (a, b) in

- a)  $R^2$ ?
- **b**)  $R^3$ ?

c)  $R^*$ ?

**25.** Use Algorithm 1 to find the transitive closures of these relations on {1, 2, 3, 4}.

- a)  $\{(1, 2), (2, 1), (2, 3), (3, 4), (4, 1)\}$
- **b**) {(2, 1), (2, 3), (3, 1), (3, 4), (4, 1), (4, 3)}
- **c)** {(1, 2), (1, 3), (1, 4), (2, 3), (2, 4), (3, 4)}
- **d**) {(1, 1), (1, 4), (2, 1), (2, 3), (3, 1), (3, 2), (3, 4), (4, 2)}

Note: avoid symmetry to become irreflexive  $a \neq b$ 

## **Equivalence Relations**

A relation on a set A is called an *equivalence relation* if it is reflexive, symmetric, and transitive.

**EXAMPLE 3** Congruence Modulo m Let m be an integer with m > 1. Show that the relation

$$R = \{(a, b) \mid a \equiv b \pmod{m}\}$$

is an equivalence relation on the set of integers.

#### **Equivalence Classes**

if R is an equivalence relation on a set A, the equivalence class of the element a is  $[a]_R = \{s \mid (a, s) \in R\}$ 

What are the equivalence classes of 0, 1, 2, and 3 for congruence modulo 4?

*Solution:* The equivalence class of 0 contains all integers a such that  $a \equiv 0 \pmod{4}$ . The integers in this class are those divisible by 4. Hence, the equivalence class of 0 for this relation is

$$[0] = {\ldots, -8, -4, 0, 4, 8, \ldots}$$


The equivalence class of 1 contains all the integers a such that  $a \equiv 1 \pmod{4}$ . The integers in this class are those that have a remainder of 1 when divided by 4. Hence, the equivalence class of 1 for this relation is

$$[1] = {\ldots, -7, -3, 1, 5, 9, \ldots}.$$

- **1.** Which of these relations on {0, 1, 2, 3} are equivalence relations? Determine the properties of an equivalence relation that the others lack.
  - a)  $\{(0,0), (1,1), (2,2), (3,3)\}$
  - **b**)  $\{(0,0), (0,2), (2,0), (2,2), (2,3), (3,2), (3,3)\}$
  - **c)**  $\{(0,0), (1,1), (1,2), (2,1), (2,2), (3,3)\}$
  - **d**) {(0, 0), (1, 1), (1, 3), (2, 2), (2, 3), (3, 1), (3, 2), (3, 3)}
  - **e**) {(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (1, 2), (2, 0), (2, 2), (3, 3)}


In Exercises 21–23 determine whether the relation with the directed graph shown is an equivalence relation.

21.




22.





23.



- a) 2?
- **b**) 3?
- **c)** 6?
- **d**) -3?

## **Partial Orderings**

A relation R on a set S is called a *partial ordering* or *partial order* if it is reflexive, antisymmetric, and transitive. A set S together with a partial ordering R is called a *partially ordered* set, or *poset*, and is denoted by (S, R). Members of S are called *elements* of the poset.

Show that the greater than or equal to relation  $(\geq)$  is a partial ordering on the set of integers.

Solution: Because  $a \ge a$  for every integer  $a, \ge$  is reflexive. If  $a \ge b$  and  $b \ge a$ , then a = b. Hence,  $\ge$  is antisymmetric. Finally,  $\ge$  is transitive because  $a \ge b$  and  $b \ge c$  imply that  $a \ge c$ . It follows that  $\ge$  is a partial ordering on the set of integers and  $(\mathbf{Z}, \ge)$  is a poset.

The divisibility relation | is a partial ordering on the set of positive integers, because it is reflexive, antisymmetric, and transitive, as was shown in Section 9.1. We see that  $(\mathbf{Z}^+, |)$  is a poset. Recall that  $(\mathbf{Z}^+$  denotes the set of positive integers.)

### Lexicographic Order

The lexicographic ordering  $\leq$  on  $A_1 \times A_2$ 

$$(a_1, a_2) \prec (b_1, b_2)$$
, either if  $a_1 \prec_1 b_1$  or if both  $a_1 = b_1$  and  $a_2 \prec_2 b_2$ .

Determine whether (3, 5) < (4, 8), whether (3, 8) < (4, 5), and whether (4, 9) < (4, 11) in the poset  $(\mathbf{Z} \times \mathbf{Z}, \leq)$ , where  $\leq$  is the lexicographic ordering constructed from the usual  $\leq$  relation on  $\mathbf{Z}$ .

Solution: Because 3 < 4, it follows that (3, 5) < (4, 8) and that (3, 8) < (4, 5). We have (4, 9) < (4, 11), because the first entries of (4, 9) and (4, 11) are the same but 9 < 11.

elexicographic ordering of strings.  $a_1 a_2 \dots a_m$  and  $b_1 b_2 \dots b_n$  $t = \min(m, n)$ 

string  $a_1 a_2 ... a_m$  is less than  $b_1 b_2 ... b_n$  if and only if  $(a_1, a_2, ..., a_t) < (b_1, b_2, ..., b_t)$ , or  $(a_1, a_2, ..., a_t) = (b_1, b_2, ..., b_t)$  and m < n,

discreet ≺ discrete, discreet ≺ discreetness discrete ≺ discretion, discrete ≺ discreti

- 1. Which of these relations on {0, 1, 2, 3} are partial orderings? Determine the properties of a partial ordering that the others lack.
  - **a)** {(0, 0), (1, 1), (2, 2), (3, 3)}
  - **b**) {(0, 0), (1, 1), (2, 0), (2, 2), (2, 3), (3, 2), (3, 3)}
  - **c**) {(0, 0), (1, 1), (1, 2), (2, 2), (3, 3)}
  - **d**) {(0, 0), (1, 1), (1, 2), (1, 3), (2, 2), (2, 3), (3, 3)}
  - **e**) {(0, 0), (0, 1), (0, 2), (1, 0), (1, 1), (1, 2), (2, 0), (2, 2), (3, 3)
- **5.** Which of these are posets?
  - a)  $(\mathbf{Z}, =)$  b)  $(\mathbf{Z}, \neq)$  c)  $(\mathbf{Z}, \geq)$

- d) (**Z**, //)
- 7. Determine whether the relations represented by these zero-one matrices are partial orders.

  - a)
      $\begin{vmatrix}
     1 & 1 & 1 \\
     1 & 1 & 0 \\
     0 & 0 & 1
     \end{vmatrix}$  b)
      $\begin{vmatrix}
     1 & 1 & 1 \\
     0 & 1 & 0 \\
     0 & 0 & 1
     \end{vmatrix}$
  - $\mathbf{c}) \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$

- **16.** Let  $S = \{1, 2, 3, 4\}$ . With respect to the lexicographic order based on the usual less than elation,
  - a) find all pairs in  $S \times S$  less than (2, 3).
  - **b**) find all pairs in  $S \times S$  greater than (3, 1).
- 18. Find the lexicographic ordering of these strings of lowercase English letters:
  - a) quack, quick, quicksilver, quicksand, quacking
  - b) open, opener, opera, operand, opened
  - c) zoo, zero, zoom, zoology, zoological
- 19. Find the lexicographic ordering of the bit strings 0, 01, 11, 001, 010, 011, 0001, and 0101 based on the ordering 0 < 1.